

**IN THE UNITED STATES PATENT
AND TRADEMARK OFFICE**

Appln No. : 10/600,266
Confirm. No.: 7488
Applicant(s): Fumitoshi ASAI et al
Filed : June 20, 2003

For : MEDICINAL COMPOSITIONS
CONTAINING ASPIRIN

Art Unit : 1614

Examiner : Brian Yong S. Kown

Docket No. : 03337C/HG

Customer No.: 01933

SUPPLEMENTAL AMENDMENT

COMMISSIONER FOR PATENTS
P.O. Box 1450
Alexandria, VA 22313-1450

S I R :

This is to supplement the amendment filed October 16, 2007
in which a Declaration of Dr. Fumitoshi Asai was attached.
The Declaration was a Declaration Under 37 CFR 1.131 to
predate a reference. To complete the record there is
enclosed herewith a substantially identical Declaration
which is signed by the other three co-inventors.

Frishauf, Holtz, Goodman
& Chick, P.C.
220 Fifth Ave., 16th Floor
New York, NY 10001-7708
Tel. No. (212) 319-4900
Fax No.: (212) 319-5101
MJC/sg
Enc. Declaration with attachments

CERTIFICATE OF FACSIMILE
TRANSMISSION TO NO.1-571-273-8300
TOTAL PAGES: 19

RECEIVED
CENTRAL FAX CENTER

OCT 22 2007

I hereby certify that this
paper is being facsimile
transmitted to the
Patents and Trademarks Office
on the date noted below.

Attorney: MARSHALL S. CHICK

Dated: October 22, 2007

In the event that this Paper
is late filed, and the necessary
petition for extension of time is
not filed concurrently herewith,
please consider this as a Petition
for the requisite extension of time,
and to the extent not tendered by
payment attached hereto,
authorization
to charge the extension fee, or any
other fee required in connection with
this Paper, to Account No. 06-1378.

Respectfully submitted,

MARSHALL S. CHICK
Reg. No. 26,853

RECEIVED
CENTRAL FAX CENTER

OCT 22 2007

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Appl. No. : 10/600,266
Confirm. No.: 7488
Applicant(s): Fumitoshi ASAI et al
Filed : June 20, 2003
Art Unit : 1614
Examiner : Brian Yong S. Kwon
Docket No. : 03337C/EG
For : MEDICINAL COMPOSITIONS
CONTAINING ASPIRIN
Customer No.: 01933

DECLARATION UNDER 37 CFR 1.131

The below named declarants hereby declare the following:

1. They are each a co-inventor of the invention described and claimed in the above-identified application.
2. Attached hereto are copies of notebook records documenting experiments done by us (the inventors) or under our supervision and control, showing a reduction to practice of the claimed invention. The code "CS 747" which appears throughout the notebook pages is our internal code for the compound identified as "Compound A" in Table I of the specification of our patent application. The dates on the copies have been blacked out. Translations of these documents are also attached. The acts described in these documents occurred prior to November 3, 1998.

We hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001, of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Date: 10/22/2007

Atsuhiko SUGIMACHI
Atsuhiko SUGIMACHI


Date: 10/22/2007

Takatoshi OGAWA
Takatoshi OGAWA

Date: 10/19/2007

Teruhiko INOUE
Teruhiko INOUE

Attachment: Notebook records and
English translation thereof

Page 145


CS 747
+ Combination Experiment
Aspirin

Dose	CS 747	0.3 mg/kg (4hr)	about 40 mg
		1 mg/kg (4hr)	about 20

) Dose in which both bleeding time and aggregation were measured:

Aspirin 10 + CS 747 0.6 2hr

First, experiment with a central focus on Aspirin 10 + CS 747 0.6

Another group 0.3 or 1

Conduct 0.3 because 1 seems to work too much.

Page 146

Arterio-venous Shunt Thrombosis Model in Rats
(Examination of effect by combination of CS-747 with aspirin)

[Object]

Examine the effect by combination of CS-747 with aspirin using Arterio-venous shunt thrombosis model in rats.

[Experimental Term]

Thirty six rats received on [REDACTED] are used.

[Animals]

Seven-week-old male SD rats (Japan SLC) are purchased and used for the experiment after preliminary breeding for about a week. The experiment is conducted as 6 rats per group.

[Test agents]

CS-747 (synthesized by Ube Industrials Ltd., Lot No. 16) and aspirin (Sigma, A-5376, Lot No. 46H1053, received on [REDACTED]) are used. The test agents are dissolved or suspended in a 5% Arabic gum (Sigma, Lot No. 73H0705, opened on [REDACTED]) solution and administered orally in volume of 1 ml/kg two hours before starting arterio-venous shunt. Administered group are (A) vehicle, (B) aspirin 10 mg/kg, (C) CS-747 0.3 mg/kg, (D) CS-747 0.6 mg/kg, (E) aspirin 10 mg/kg + CS-747 0.3 mg/kg, and (F) aspirin 10 mg/kg + CS-747 0.6 mg/kg.

[Methods]

(1) For the experiment, the method by Umetsu et al. (Thromb. Haemost. 39, 74-83, 1978) is partly modified.

(2) The shunt tube for arterio-venous shunt is prepared as follows; both sides of a medical silicon tube of 12 cm length (inner diameter: 1.5 mm, outer diameter: 2.5 mm, KANEKA Medix Co., Ltd) are connected each to a polyethylene tube of 7 cm length (inner diameter: 0.5 mm, outer diameter: 1.0 mm, Natsume Seisakusho Co., Ltd.)

covered with silicon via a medical silicon tube of 0.7 cm length (inner diameter: 1.0 mm, outer diameter: 1.5 mm, KANEKA Medix Co., Ltd.) as connector. At the connection,

Page 147
[REDACTED]

surgical adhesive (Aronalpha A, Sankyo) is used for preventing blood leak. In addition, a silk thread (size 3-0, Niccho Kogyo) of 10 cm length is placed in the tube of 12 cm length.

(3) Vehicle (5% Arabic gum solution) or test agents are administered orally in a volume of 1 ml/kg 2 hours before starting arterio-venous shunt. 6 rats per group are used.

(4) The above tube prepared in advance is filled with heparin solution (Japanese Pharmacopoeia Heparin Sodium Injection, Fuso Pharmaceutical Industries, Ltd., Lot No. 97H28A, received on [REDACTED]) diluted with normal saline (Otsuka) resulting in 30 unit/kg.

(5) The rat is anesthetized with an intraperitoneal injection of 1 ml/kg (40 mg/kg) of pentobarbital solution (Nembutal R, Abbott, Lot No. 20-975-27) diluted with normal saline resulting in 40 mg/ml. After it is fixed to turning up, the jugular vein is exposed and one side of the shunt tube (in which the silk thread is not adhered) is cannulated. Subsequently, to the carotid artery where bloodstream is shut using clamp, the other side of the tube is cannulated to make the arterio-venous shunt.

(6) After removing the clamp and allowing blood to circulate for 30 minutes, the thrombus adsorbed on the silk thread is weighed. The thrombus weight was calculated by subtracting of the weight of the thread (6.5 mg) from the measured weight.

Files were stored at AV-shunt (3) (F00515 data)

[REDACTED] A. Sugidachi

Page 148

AV shunt Thrombosis Model in Rats
(Combination of CS 747 with aspirin)

Protocol: P. 146, 147

Reagent and so on: p. 110, p. 138

Cage number: GraC3-01-04

Rats

SD male (Japan SLC)

Sex, system: male SD

wks, Receipt

Year-round old: 7 weeks

Receipt number: 034163

Body weight:

Sugidachi

Manufacture name: Japan SLC

Receive date: [REDACTED]

Number of rats: 36

Experimenter: Atsuhiko Sugidachi

Housing term: [REDACTED] to [REDACTED]

Receipt number: 034163

5% Arabic gum soln.

67.7mg aspirin

50 mg/ml = 1011.7 mg / 20.234 ml dH₂O

16.2 mg CS 747

Aspirin

10 mg/ml = 67.7 mg / 5.77 ml 5% Arabic gum soln.

1011.7 mg Arabic gum

CS 747

16.2 mg / 5.4 ml = 3 mg/ml

3 mg/ml soln. 1 ml + 5% Arabic gum soln. 2 ml = 1 mg/ml

1 mg/ml soln. 1 ml + 5% Arabic gum soln. 2.33 ml = 0.3 mg/ml

1 mg/ml soln. 1.5 ml + 5% Arabic gum soln. 1 ml = 0.6 mg/ml

10.7 mg CS 747

Further prepared because of insufficient (spilled)

5% Arabic gum 1084.7 mg / 21.69 ml dH₂O = 50 mg/ml

1084.7 mg Arabic gum

CS 747 10.7 mg / 10.7 ml Arabic gum soln. = 1 mg/ml

1 mg/ml soln. 1.5 ml + 5% Arabic gum soln. 3.5 ml = 0.3 mg/ml

1 mg/ml soln. 3 ml + 5% Arabic gum soln. 2 ml = 0.6 mg/ml

Page 149

	B. W. (g)	Treatment	Measured value (mg)	Thrombus (mg)
#1	253	Vehicle	61.3	54.8
2	252	Aspirin 10	51.3	44.8
3	262	CS 747 0.3	58.3	51.8
4	267	CS 747 0.6	43.1	36.6
5	256	Aspirin 10 + CS 747 0.3	39.6	33.1
6	271	Aspirin 10 + CS 747 0.6	23.2	16.7
7	250	V	63.2	56.7
8	246	A 10	58.4	51.9
9	258	747 0.3	51.8	45.3
10	269	747 0.6	53.1	46.6
11	268	A 10 + 747 0.3	30.5	24.0
12	244	A 10 + 747 0.6	41.3	34.8
13	247	V	56.1	49.6
14	262	A 10	48.6	42.1
15	256	747 0.3	52.2	45.7
16	267	747 0.6	46.3	39.8
17	268	A 10 + 747 0.3	42.2	36.7
18	242	A 10 + 747 0.6	21.6	15.1

V = Vehicle

A = Aspirin

747 = CS 747

heparin

1000 u/ml soln. 1 ml + saline 9 ml = 100 unit/ml

100 u/ml soln. 3 ml + saline 7 ml = 30 unit/ml

#1 252.9 g
251.7 g
262.3 g
267.3 g
256.4 g
271.1 g

#7 250.0 g
245.8 g
257.8 g
269.0 g
268.3 g
244.4 g

#13 247.0 g
262.3 g
266.2 g
267.3 g
267.6 g
242.1 g

Page 150

[REDACTED]

#1	0.0613 g	#7	0.0632 g	#13	0.0561 g
#2	0.0513 g	#8	0.0584g	#14	48.6 mg
#3	0.0583 g	#9	0.0518 g	#15	52.2 mg
#4	0.0431 g	#10	0.0531 g	#16	0.0463 g
#5	0.0396 g	#11	0.0305 g	#17	0.0422 g
#6	0.0232 g	#12	41.3 mg	#18	0.0216 g

[REDACTED] A. Sugidachi

Page 151

AV Shunt Thrombosis Model in Rats
(Aspirin + CS 747)

Protocol: p. 146, 147

Reagents and so on: p. 110, 138

Cage number: GraC3-01-01

Rat

6

SD-male (7 wks, Receipt)

Sex, system: male SD

36+2

Japan SLC

Year-round old: 7 weeks

Receipt

Body weight:

Sugidachi

Manufacture name: Japan SLC

Receipt number: 034163

Receive date: [REDACTED]

Number of rats: 36

Experimenter: Atsuhiko Sugidachi

Housing term: [REDACTED] to [REDACTED]

Receipt number: 034163

Additional 2 rats

↓
euthanasia using CO₂ gas

[REDACTED]	B. W.	72.0 mg
#1	271.0 g	11.3 mg
	277.6 g	
	272.8 g	
	275.0 g	
	245.5 g	
#6	251.6 g	1224.1 mg
#10	248.2 g	
#8	263.9 g	
#9	253.3 g	
#7	281.1 g	
#11	263.4 g	
#12	263.4 g	
		5% Arabic gum = 50 mg/ml
		= 1224.1 / 24.48 ml dH ₂ O
#13	271.3 g	
	265.3 g	
	266.0 g	
	255.5 g	
	245.6 g	
#15	272.4 g	

Aspirin

10 mg/ml = 72.0 / 7.2 ml 5% Arabic gum soln.

Page 152

CS 747 11.3 mg / 11.3 ml 5% Arabic gum soln. = 1 mg/ml
 1 mg/ml soln. 1.5 ml + 5% Arabic gum soln. 1ml = 0.6 mg/ml
 1 mg/ml soln. 1 ml + 5% Arabic gum soln. 2.33 ml = 0.3 mg/ml

heparin 1000 unit/ml soln. (origine) 1 ml + saline 9 ml = 100 unit/ml
 100 unit/ml soln. 3 ml + saline 7 ml = 30 unit/ml

	B. W. (g)	Treatment	Measured value (mg)	Thrombus (mg)
#1	271	V	55.6	49.1
2	278	A <u>10</u>	52.9	46.4
3	273	747 <u>0.3</u>	43.9	37.4
4	275	747 <u>0.6</u>	41.0	34.5
5	246	A <u>10</u> + 747 <u>0.3</u>	24.8	18.1
6	252	A <u>10</u> + 747 <u>0.6</u>	19.7	13.2
7	281	V	59.0	52.5
8	264	A <u>10</u>	63.2	56.7
9	253	747 <u>0.3</u>	47.1	40.6
10	248	747 <u>0.6</u>	38.5	32.0
11	263	A <u>10</u> + 747 <u>0.3</u>	36.7	30.2
12	263	A <u>10</u> + 747 <u>0.6</u>	35.6	29.1
13	271	V	57.8	51.3
14	265	A <u>10</u>	44.3	37.8
15	266	747 <u>0.3</u>	46.4	39.9
16	256	747 <u>0.6</u>	42.2	35.7
17	246	A <u>10</u> + 747 <u>0.3</u>	48.3	41.8
18	272	A <u>10</u> + 747 <u>0.6</u>	36.6	30.1

V = Vehicle

A = Aspirin

747 = CS 747

Page 153
[REDACTED]


#1	0.0556 g	#7	0.0590 g	#13	0.0578 g
#2	0.0529 g	#8	0.0632 g	#14	44.3 mg
#3	0.0439 g	#9	0.0471 g	#15	0.0464 g
#4	0.0410 g	#10	0.0385 g	#16	42.2 mg
#5	0.0246 g	#11	0.0367 g	#17	48.3 mg
#6	0.0197 g	#12	0.0356 g	#18	36.6 mg

[REDACTED] A. Sugidachi

[Hirose] [REDACTED]

Page 154

Arterio-venous shunt thrombosis model in rats



Aspirin + CS 747 Summary

Vehicle	52.3 ± 1.2
Aspirin 10	46.6 ± 2.8
CS 747 0.3	43.5 ± 2.1
CS 747 0.6	37.5 ± 2.1
Aspirin 10 + CS 747 0.3	30.5 ± 3.5
Aspirin 10 + CS 747 0.6	23.2 ± 3.8

A. Sugidachi

145

CS 747

Aspirin

1/14 Exp. 行なう。

Dose CS 747 0.3 mg/kg (4hr) で 17 40 mg.
1 mg/kg (4hr) で 17 20

出血時間, 凝集 なども行う。 12

Aspirin ⑩ + CS 747 ⑥ 2hr

33 Aspirin ⑩ + CS 747 ⑥ を中心として行う。

ES 121 ⑥ - ⑦ - ⑦は 20分おきで ⑥ と行う。

[1
ラ
る。

[3

[10
雄
い[課
CS-
46H
No.
合で
(B)
aspir
する[方
(1) 3
して(2) 3
ディ
の両
ーブ
で0.1
カネ

147

ラット

ロンアルファA、三共)で接種する。また12 cmのチューブ内に10 cmの絹糸
(3-0、日腸工業)を設置する。

(3) 動脈シャント開始2時間前に、vehicle (5%アラビアゴム溶液) または薬物
を1 ml/kgの割合で経口投与する。1群6匹の実験を行う。

(4) あらかじめ作成しておいた上記チューブに30 unit/kgとなるように、生理食
塩液 (大塚) で希釈したヘパリン溶液 (日本薬局方ヘパリンナトリウム注射液、
扶桑薬品工業、Lot No. 97H28A、XXXXXXXXXX入荷) を満たす。

(5) 生理食塩液で40 mg/mlに希釈したベントバルビタール溶液 (ネンブタール®、
Abbott、Lot No. 20-975-Z7) を1 ml/kg腹腔内投与 (40 mg/kg) してラットを麻酔
させる。仰臥位に固定させた後、頸静脈を露出させ、シャント用チューブの片
側 (糸の付いていない方) をカニューレーションする。続いて、クレンメで血流
を遮断しておいた頸動脈に、チューブの反対端をカニューレーションし、動脈
シャントを形成する。

(6) クレンメをはずしてシャントに30分間血液を循環させた後、絹糸に付着し
た血栓の湿重量を測定する。測定重量から糸の重量 (6.5 mg) を差し引き血栓
重量を求める。

7°

ケージ
性別、
周3 薬
入
実
飼育
受付57.
16.

1011.

ファイル AV-shunt (3) 1 = 保存
(F005157-9)

10.7

1084.7

A. Sigita

149

	B.W. (g)	Treatment	測定値 (ms)	Thrombus (mg)	
#1	253	Vehicle	61.3	54.8	
2	252	Aspirin (10)	51.3	44.8	#1 B.
3	262	CS 1747 (0.3)	58.3	51.8	
4	267	CS 1747 (0.6)	43.1	36.6	
5	256	Aspirin (10) + CS 1747 (0.3)	39.6	33.1	
6	271	Aspirin (10) + CS 1747 (0.6)	23.2	16.7	
7	250	V	63.2	56.7	#2 B.
8	246	A (10)	58.4	51.9	
9	258	1747 (0.3)	51.8	45.3	
10	269	1747 (0.6)	53.1	46.6	
11	268	A (10) + 1747 (0.3)	30.5	24.0	
12	244	A (10) + 1747 (0.6)	41.3	34.8	#3 B.
13	247	V	56.1	49.6	
14	262	A (10)	48.6	42.1	
15	256	1747 (0.3)	52.2	45.7	
16	267	1747 (0.6)	46.3	39.8	#4 B.
17	268	A (10) + 1747 (0.3)	42.2	35.7	
18	242	A (10) + 1747 (0.6)	21.6	15.1	

V = Vehicle

A = Aspirin

1747 = CS 1747

ハロニン

1000 u/ml soln. 1ml + saline 9ml = 100 unit/ml

100 u/ml soln. 3ml + saline 7ml = 30 unit/ml

#1 252.9g
251.7g
262.3g
267.3g
256.4g
271.1g

#7 250.0g
245.8g
257.8g
269.0g
266.3g
244.4g

#13 247.8g
262.3g
256.2g
267.3g
267.6g
242.1g

01-10-12;17:30

Daiichi Sanyo IP

GOODMAN

0336805623

36/

151

ラット AV shut 血圧モデル
(アスピリン + CS744)

7.0 コール

p146, 147

計測

p110, 118

CS74

ケージ番号: 67C3-01-01

性別、系統: ♂ SD
周年令: 7週

体重: 日本SLC

飼育者名: 杉立 取寛

飼育期間: 034163

受付番号: 034163

6
36 + 2

匹数: 36

ラット

SD 含 (7wks 入荷)

日本SLC

入荷

杉立

受付番号 034163

7/10

B.W.

#1 271

2 278

3 273

4 275

5 246

6 252

7 281

8 264

9 253

10 248

11 263

12 263

13 271

14 265

15 266

16 256

17 246

18 272

おまけ2匹

↓

CO2 gas 2匹

± 5%

B.W.

+1 271.0g
277.6g
272.8g
275.0g
245.5g
+2 251.6g

#10 248.2g
#2 263.9g
#9 253.3g
#7 281.1g
#11 263.4g
#17 263.4g

#13 271.3g
265.3g
266.0g
258.5g
245.6g
#18 272.4g

72.0 mg
11.3 mg

1224.1 mg

5% アスピリン = 50 mg/ml

= 1224.1 / 24.48 ml d H2O

Aspirin

10 mg/ml = 72.0 / 7.2 ml 5% アスピリン sol'n.

153

			Thrombus weight (mg)
#1	0.0556 g	#7	0.0590 g
		#13	0.0578 g
#2	0.0529 g	#8	0.0632 g
		#14	44.3 mg
#3	0.0439 g	#9	0.0471 g
		#15	0.0464 g
#4	0.0410 g	#10	0.0385 g
		#16	42.2 mg
#5	0.0246 g	#11	0.0367 g
		#17	48.3 mg
#6	0.0197 g	#12	0.0356 g
		#18	36.6 mg

Aspirin + C

Vel
A1
C
CS
A1
A1

A. Sugawara

